

Preventing
Teen
Cough
Medicine
Abuse



A PARENT'S GUIDE

You already know about the dangers of illegal street drugs like marijuana, cocaine, and methamphetamine. But did you know that some teens are abusing legal products, like cough medicine, to get high?

This guide will provide the answers to many questions about cough medicine abuse, give helpful tips for preventing this and other types of drug abuse, and offer other resources for more information.

What is cough medicine abuse?

Cough medicine abuse is taking extremely large doses of cough medicine to get high. The “high” is caused by taking a large amount of dextromethorphan, which is often abbreviated DXM, a common active ingredient found in many cough medications. This sort of abuse—whether it’s called cough medicine abuse, or dextromethorphan or DXM abuse—can be dangerous.

What is dextromethorphan?

Dextromethorphan is a safe and effective active ingredient found in many nonprescription, over-the-counter (OTC) cough medicines, including syrups, tablets, and gel caps. When used according to medicine label directions, the ingredient dextromethorphan produces few side effects and has a long history of safety. When abused in large amounts, it can produce a “high” feeling as well as a number of dangerous side effects.

What are the effects of cough medicine abuse?

The effects of abusing cough medicines containing dextromethorphan vary with the amount taken. Common effects when taken in large amounts include confusion, dizziness, double or blurred vision, slurred speech, loss of physical coordination, abdominal pain, nausea and vomiting, rapid heart beat, drowsiness, numbness of fingers and toes, and disorientation.

DXM abusers describe different “plateaus” ranging from mild distortions of color and sound to visual hallucinations, “out-of-body” dissociative sensations, and loss of motor control. Side effects can be worsened if the dextromethorphan-containing cough medicine being abused also contains other ingredients to treat more than just coughs.

Cough medicine also is sometimes abused in combination with other medications, alcohol, and illegal drugs, which can increase the dangerous side effects.

What cough medicines contain dextromethorphan, or DXM?

There are over 100 OTC medicines that contain DXM, either as the only active ingredient or in combination with other active ingredients. Examples include Alka-Seltzer Plus Cold & Cough Formula, Coricidin HBP Cough and Cold, Dimetapp® DM, Mucinex® DM tablets, PediaCare cough medicines, certain Robitussin® cough products, Sudafed cough products, TheraFlu Cough products, Triaminic cough products, Tylenol Cough and Tylenol Cold products, Vicks 44 Cough Relief products, and certain Vicks DayQuil and NyQuil LiquiCaps. A number of store brands contain dextromethorphan, as well.

To know if a product contains DXM, look for “dextromethorphan” in the *active ingredient* section of the OTC Drug Facts label. You also may look for an educational icon on the packaging of most OTC cough medicines containing dextromethorphan.

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

This icon serves as a mini public service announcement for parents, making them aware of cough medicine abuse among teens, and points them to StopMedicineAbuse.org where they can find resources to talk to their teens about the issue.

What are slang terms for dextromethorphan?

The most common terms include Dex, DXM, Robo, Skittles, Syrup, Triple-C, and Tussin. Terms for using dextromethorphan include: Robo-ing, Robo-tripping, and Skittling, among others. Slang terms for dextromethorphan vary by product and region. Adults should be familiar with all of them.

How common is cough medicine abuse?

Research shows that one out of 10 teens, or over two million teens, from across the country and of all backgrounds, has abused cough medicine to get high.

Where are teens finding information about cough medicine abuse?

Often, these teens find information about how to abuse cough medicine on the Internet. A number of web sites promote the abuse of cough medicines containing DXM. Some of these sites even recommend how much to take; suggest other drugs to combine with DXM; instruct how to extract DXM from cough medicines; promote drug abuse in general; or even sell a raw, unfinished form of DXM for snorting. Be aware of what your teen does on the Internet, the web sites he or she visits, and the amount of time he or she is logged on.

The abuse of cough medicine also now can be seen in some current music, movies, and fashion.

What Parents Can Do to Prevent Cough Medicine Abuse

Parents can make a positive difference in their teens' lives, and research shows that parents do influence their teens' decisions about whether to take drugs or not. To prevent DXM abuse specifically, the best advice is to educate yourself, communicate with your teens, and safeguard your medications.

Educate Yourself

- Educate yourself about cough medicine abuse and share this information with others who are in contact with your teen, such as school administrators, coaches, and counselors.
- Know the signs of cough medicine abuse and what to watch out for.

Communicate with Your Teens

- Be clear that you do not want your teen taking medicine without your knowledge.
- Teach your teens and younger children to respect medicines. Medicines are important tools in healthcare, but they must be used according to directions.
- Make sure your teen understands that abusing cough medicine—just as abusing illegal drugs—can be very dangerous.

Safeguard Medications

- Know what medicines are in your home and pay attention to quantities.
- Keep all medications out of reach and out of easily accessible places like medicine cabinets.
- If your child needs medications during school hours, speak with school officials about medicine policies in the school.

More Tips for Raising Drug-Free Teens

Monitoring is an effective way you can help your teen stay drug-free, and an important thing to do even if you don't suspect your teen is using drugs. Monitoring means asking young people questions about where they're going, what they're doing, and with whom they're spending time, as well as

keeping tabs on their Internet use by using web browser tools and software designed to block certain sites. Put some of these tips to use, and your kids will benefit.

Talk with Your Teen

Teens who learn a lot about the dangers of drugs from their parents are up to half as likely to abuse drugs. Parents should talk often, listen regularly, and communicate clearly that they do not want their kids using drugs.

To talk credibly and effectively about the dangers of drugs, parents need to know what those dangers are. Teens know when they're being lied to, so know the facts. To learn about the various drugs teens can abuse—including cough medicine—and the risks, visit the Partnership for a Drug-Free America at drugfree.org.

Know Where Your Teen Is

It's important to know where your teen is and what he or she is doing. Research has shown that children without adult supervision are at significantly greater risk of truancy from school, stress, receiving poor grades, risk-taking behavior, and substance abuse.

Introduce Your Teen to Adult Role Models

Find out what adult-supervised activities—like clubs or after-school sports—interest your teen and help get him or her involved. Connect with other adults who can help teens avoid the dangers of drugs, and reinforce the benefits of healthy, drug-free living.

Know Your Teen's Friends

Research from the Partnership for a Drug-Free America reports that more than half of teens say they have close friends who get high regularly. Parents need to know if these are the friends with whom their children are spending time.

Recognize Signs Your Teen Is Using Drugs

Parents don't always recognize their kids might be using drugs. While it can be hard to know, there are some general warning signs you can watch for. The fact is, any teen could be using drugs, so stay alert. As a general rule, changes that are sudden or extreme may be a warning sign.

Signs your teen could be using drugs include:

- Change in friends
- Change in eating or sleeping patterns
- Changes in physical appearance and hygiene
- Declining grades
- Loss of interest in hobbies or favorite activities
- Hostile and uncooperative attitude
- Unexplained disappearance of household money
- Visits to pro-drug web sites
- Empty drug or medicine containers or drug paraphernalia
- Unusual chemical or medicinal smells on your child or in his or her room

Helping a Teen Who Is Using Drugs

The goal is to prevent a drug abuse problem in the first place. If you fear, however, that your teen may have a problem, sit down with your child for an open discussion about alcohol and drug use. Openly voice your suspicions but avoid direct accusations. Do not have this conversation when your teen is under the influence of alcohol or other drugs, and make sure you sound calm and rational. Ask your teen what has been going on in his or her life. Discuss ways to avoid using alcohol and other drugs in the future. If you need help during this conversation, ask another family member, your child's guidance counselor, or a physician.

Be firm and enforce whatever discipline you've laid out in the past for breaking house rules. You also should discuss ways your teen can regain your lost trust, such as calling in, spending evenings at home, or improving grades.

Resources for More Information

If you think your teen needs professional help, your doctor, hospital, or school nurse or counselor may be able to help. Or you can call 1.800.662.HELP or visit <http://findtreatment.samhsa.gov>, the treatment locator hotline and web site of the U.S. Substance Abuse and Mental Health Services Administration.

If you suspect a poisoning,
call your local poison control center at
1.800.222.1222.

To learn more about drug prevention,
visit the Partnership for a Drug-Free America at
drugfree.org.

For additional copies of this brochure, visit
StopMedicineAbuse.org.



**CHPA EDUCATIONAL
FOUNDATION**



Mixed Sources

Product group from well-managed
forests, controlled sources and
recycled wood or fiber
www.fsc.org Cert no.
© 1996 Forest Stewardship Council